



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Diffuse Lung Disease

TYPE: Abstract

TOPIC: Diffuse Lung Disease

PREDICTORS OF POST-COVID-19 PULMONARY FIBROSIS: A CASE SERIES

R.-E. CÎRJALIU^{*1} A.-P. FILDAN² AND D. TOFOLEAN² ¹OVIDIUS UNIVERSITY OF CONSTANTA, 5TH DEPARTMENT OF CLINICAL INTERNAL DISCIPLINES, CONSTANTA, ROMANIA ²OVIDIUS UNIVERSITY OF CONSTANTA/FACULTY OF MEDICINE, 5TH DEPARTMENT OF CLINICAL INTERNAL DISCIPLINES, CONSTANTA, ROMANIA

PURPOSE: The characteristics of pulmonary fibrosis in patients with COVID-19 (coronavirus disease 2019) is not well known yet. Pulmonary fibrosis determines non-reversible lung dysfunction. The aim of this study is to identify the risk factors for pulmonary fibrosis in COVID-19 patients so that we can identify early the patients with high risk and administrate proper therapy.

METHODS: From September 2020 to September 2021, using follow-up CT (computed tomography) scans we identified 31 patients with lung fibrosis, who previously had COVID-19.

RESULTS: From the population analyzed we identified as a risk factor advanced age, 67% of the patients being over 60 years old. We also found that 77% of the patients were smokers. The incidence of pulmonary fibrosis was higher in patients with severe form of COVID-19 (77%), than the ones who presented moderate form. The most frequent comorbidities found were arterial hypertension (61%) and obesity (51%). The laboratory findings associated with pulmonary fibrosis were leukocytosis, increased LDH, and inflammatory syndrome. The majority of the patients presented lung function abnormalities, such as restrictive syndrome (61%) impaired diffusion capacity (58%). Most of the patients also presented reduced 6-minute-walk test distance (80%).

CONCLUSIONS: Although there is not a specific cause know yet for post-COVID-19 lung fibrosis, there are some predicting factors, such as cigarette smoking, arterial hypertension, obesity, severe form of COVID-19, leukocytosis, inflammatory syndrome, and increased LDH.

CLINICAL IMPLICATIONS: Early detection of patients with high risk to develop post-COVID-19 pulmonary fibrosis may help the prevention and management of this severe complication.

DISCLOSURE: Nothing to declare.

KEYWORD: post-COVID-19 lung fibrosis

DOI: <http://dx.doi.org/10.1016/j.chest.2021.12.288>

Copyright © 2021 American College of Chest Physicians. Published by Elsevier Inc. All rights reserved.